

## Abstract of the Disclosure

The present invention relates to new genomic nucleotide sequences and amino acid sequences corresponding to the coding region of these genomes. The invention relates to new HCV types and subtypes sequences which are different from the known HCV types and subtypes. More particularly, the present invention relates to new HCV type 7 sequences, new HCV type 9 sequences, new HCV type 10 and new HCV type 11 sequences. Also, the present invention relates to new HCV type 1 sequences of subtypes 1d, 1e, 1f and 1g; new HCV type 2 sequences of subtypes 2e, 2f, 2g, 2h, 2l, 2k and 2i; new HCV type 3 sequences of subtype 3g, new HCV type 4 sequences of subtypes 4k, 4l and 4m; a process for preparing them, and their use for diagnosis, prophylaxis and therapy. More particularly, the present invention provides new type-specific sequences of the Core, the E1 and the NS5 regions of new HCV types 7, 9, 10 and 11, as well as of new variants (subtypes) of HCV types 1, 2, 3 and 4. These new HCV sequences are useful to diagnose the presence of HCV type 1, and/or type 2, and/or type 3, and/or type 4, and/or type 7, and/or 9, and/or type 10, and/or type 11 genotypes or serotypes in a biological sample. Moreover, the availability of these new type-specific sequences can increase the overall sensitivity of HCV detection and should also prove to be useful for prophylactic and therapeutic purposes.